Economics 620, Lecture 8a: Asymptotics II

Nicholas M. Kiefer

Cornell University

.∋...>

Suppose $\bar{X}_n - \mu \to 0$ in probability. (What can be said about the distribution of $\bar{X}_n - \mu$?) In order to get distribution theory, we need to norm the random variable; we usually look at $n^{1/2}(\bar{X}_n - \mu)$. Note that the random variable sequence $\{n(\bar{X}_n - \mu), n \ge 1\}$ does not converge in probability. (Why not?).

We might be able to make probability statements like

$$\lim_{n\to\infty} P(n^{1/2}(\bar{X}_n - \mu) < z) = F(z)$$

for some distribution F.

Then we could use F as an *approximate* distribution for $n^{1/2}(\bar{X}_n - \mu)$. This implies an *approximate* distribution for \bar{X}_n .

It is often easier to work with $Y_n = n^{1/2} (\bar{X}_n - \mu) / \sigma$.

・ロン ・聞と ・ヨン ・ヨン … ヨ

Definition: The moment generating function for the rv X (or the distribution f) is

$$m_X(t) = E(e^{tX}) = \int_{-\infty}^{\infty} e^{tx} f(x) dx.$$

The name comes from the fact that

$$\frac{d^r m}{dt^r} = \int_{-\infty}^{\infty} x^r e^{tx} f(x) dx = E(X^r)$$

when evaluated at t = 0

The subscript is dropped when unnecessary.

Note the series expansion:

$$m(t) = E(e^{tX}) = E(1 + Xt + \frac{1}{2!}(Xt)^2 + ...)$$

= $1 + \alpha_1 t + \frac{1}{2}\alpha_2 t^2 + ...$

where $\alpha_r = EX^r$ (For example: $\alpha_1 = \mu, \ \alpha_2 = \mu^2 + \sigma^2$).

Property 1: The moment generating function of $\sum_{i=1}^{n} X_i$ when X_i are independent is the product of the moment generating functions of X_i . (**Exercise**: *Prove this*.)

- * 御 > * 注 > * 注 > … 注

Property 2: Let X and Y be random variables with continuous densities f(x) and g(y). If the moment generating function of X is equal to that of Y in an interval -h < t < h, then f = g.

Example: The moment generating function for $X \sim N(0, 1)$ is

$$m(t) = E(e^{tx}) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} e^{tx} e^{-\frac{1}{2}x^2} dx$$
$$= e^{t^2/2} \left(\frac{1}{\sqrt{2\pi}}\right) \int_{-\infty}^{\infty} e^{-\frac{1}{2}(x-t)^2} dx = ?$$

The mgf does not always exist. The function $\phi_X(t) = Ee^{itX}$ always exists and is continuous and bounded.

We know
$$X_n \xrightarrow{d} X \Rightarrow \phi_{X_n}(t) \to \phi_X(t)$$
.
The converse

$$\phi_{X_n}(t) \to \phi_X(t) \; (\forall t) \Rightarrow X_n \stackrel{d}{\to} X$$

is also true

If
$$E|X| < \infty$$
 then $\phi'(0) = iEX$.
Similarly, if $EX^2 < \infty$ then $\phi''(0) = -EX^2$, etc.
CFs combine like mgfs.

3

$$\phi_X(t) = Ee^{itX} = E\cos tX + iE\sin tX$$

Thus $\phi(0) = 1, |\phi(t)| \le 1$, and $\phi(t) = \overline{\phi(-t)}$ where the bar indicates complex conjugate.

If X is symmetrically distributed, ϕ is real-valued. Some cfs: Degenerate at $\mu, \phi(t) = e^{it\mu}$

Binomial(n,p),
$$\phi(t) = (pe^{it} + 1 - p)^n$$

$$N(\mu, \sigma^2), \phi(t) = e^{it\mu - \sigma^2 t^2/2}$$

Suppose $\{X_i\}$ are iid each with cf $\phi(t)$ Then,

$$Ee^{itX} = \phi^n(t/n) = (1 + it\mu/n + o(1/n))^n$$

$$\rightarrow e^{it\mu}$$

The cf of the constant μ .

Application 2 will give a central limit theorem.

Central Limit Theorem: (CLT) (Lindberg-Levy) The distribution of $Y_n = n^{1/2}(\bar{X}_n - \mu)/\sigma$ as $n \to \infty$ is

$$\Phi(z) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{z} e^{-x^2/2} dx$$

(standard normal)

Proof: Let $\phi_{X_i-\mu}(t)$ be the characteristic function of $(X_i - \mu)$. That is

$$\phi_{X_i-\mu}(t) = 1 - \frac{\sigma^2 t^2}{2} + o(t^2)$$

where $o(t^2)$ is the remainder term such that $o(t^2)/t^2 \to 0$ as $t \to 0$.

・ロン ・聞と ・ヨン ・ヨン … ヨ

We know that

$$Y_n = \frac{\sqrt{n}(\bar{X}_n - \mu)}{\sigma} = \frac{\sum_{i=1}^n (X_i - \mu)}{\sigma \sqrt{n}},$$

Hence

$$\begin{split} \phi_{Y_n}(t) &= \left[\phi_{X_i-\mu}\left(\frac{t}{\sigma\sqrt{n}}\right)\right]^n \\ &= \left[1 - \frac{t^2}{2n} + o\left(\frac{t^2}{n}\right)\right]^n \\ &\Rightarrow \ln \phi_{Y_n}(t) = n \ln \left[1 - \frac{t^2}{2n} + o\left(\frac{1}{n}\right)\right] \\ &\approx n \ln \left[1 - \frac{t^2}{2n}\right] \end{split}$$

2

メロト メポト メヨト メヨト

$$ln\phi_{Y_n}(t) \approx n \ln \left[1 - \frac{t^2}{2n}\right]$$

 $\Rightarrow \phi_{Y_n}(t) \rightarrow e^{-t^2/2}$

as $n \to \infty$ (using $\ln(1 + x) \approx x$ for small x) which is the cf of a standard normal random variable.

Point of the Central Limit Theorem: The distribution function of \bar{X}_n for large *n* can be approximated by that of a normal with mean μ and variance σ^2/n .

1. Identical means and variances can be dropped straightforwardly. We need some restrictions on the variance sequence though. In this case, we work with

$$Y_n = \frac{\sum_{i=1}^n (X_i - \mu_i)}{\left(\sum_{i=1}^n \sigma_i^2\right)^{1/2}}.$$

2. Versions of the Central Limit Theorem with random vectors are also available. Just apply univariate theorems to all linear combinations.

3. The basic requirement is that each term in the sum should make a negligible contribution.

1. Estimation of mean μ from a sample of normal random variables: In this case, we estimate μ by \bar{X} , and the asymptotic approximation for the distribution of \bar{X} or $(\bar{X} - \mu)$ is **exact**.

2. Consider $n^{1/2}(\hat{\beta} - \beta)$ where $\hat{\beta}$ is the LS estimator.

$$n^{1/2}(\hat{\beta} - \beta) = n^{1/2}(X'X)^{-1}X'\varepsilon$$

= $[X'X/n]^{-1}n^{1/2}[X'\varepsilon/n]$

Where [X'X/n] is the sample second moment matrix of the regressors. [X'X/n] is O(1) or maybe $O_p(1)$ depending on assumptions. Its lim or plim is Q, a KxK p.d. matrix.

(本間) (本語) (本語) (二語

What about $n^{1/2}[X'\varepsilon/n] = \sqrt{n}(1/n) \sum x'_i \epsilon_i$?

This is
$$\sqrt{n}$$
 times a sample mean of $x'_i \epsilon_i$. These have $Ex'_i \epsilon_i = 0$, $Vx'_i \epsilon_i = \sigma^2 Q$ (discuss)

Under the assumption that regressors are well-behaved (i.e., contribution of any particular observation to $[X'\varepsilon/n]$ is negligible), we can apply a Central Limit Theorem and conclude that

$$n^{1/2}(\hat{\beta}-\beta) = [X'X/n]^{-1}n^{1/2}[X'\varepsilon/n] \xrightarrow{D} N(0,\sigma^2Q^{-1}).$$

Consistent with previous results?

The delta method is a "trick" for approximating the limiting distribution of a function of a statistic whose limiting distribution is known. From the CMT we know that $X_n \xrightarrow{d} X \Rightarrow g(X_n) \xrightarrow{d} g(X)$.

Suppose $X_n = \sqrt{n}(\overline{X}_n - \mu)$. What can we say about $\sqrt{n}(g(\overline{X}_n) - g(\mu))$? Expand

$$g(\overline{X}_n) \approx g(\mu) + g'(\mu)(\overline{X}_n - \mu)$$

So

$$\sqrt{n}(g(\overline{X}_n) - g(\mu)) \approx g'(\mu)\sqrt{n}(\overline{X}_n - \mu)$$

Hence if $\sqrt{n}(\overline{X}_n - \mu) \rightarrow N(0, \sigma^2)$ Then

$$\sqrt{n}(g(\overline{X}_n) - g(\mu)) \rightarrow N(0, g'(\mu)^2 \sigma^2)$$

3

- 4 週 ト - 4 ヨ ト - 4 ヨ ト - -

Suppose $\{X_i\}$ are K-variate rv's , $EX = \mu, VX = \Sigma$

Then we consider $Y = t'X - t'\mu$, univariate with EY = 0, $VY = t'\Sigma t$ and apply our CLT to conclude

$$\sqrt{n}(\overline{X_n}-\mu) \to N(0,\Sigma)$$

For the Delta method, suppose $g : \mathbb{R}^K \to \mathbb{R}^m$ and suppose $\sqrt{n}(\overline{X}_n - \mu) \to N(0, \Sigma)$

Write $g(\overline{X}_n) \approx g(\mu) + g'(\mu)(\overline{X}_n - \mu)$ where $g'(\mu)$ is the mxk matrix of derivatives and conclude

$$\sqrt{n}(g(\overline{X}_n) - g(\mu)) \rightarrow N(0, g'(\mu)\Sigma(g'(\mu))^T)$$

- 本間 と えき と えき とうき